

Existence of a metallic phase and upper bounds of the Hartree-Fock energy in the homogeneous electron gas

F. Delyon,¹ M. Duneau,¹ B. Bernu,² and M. Holzmann^{2,3}

¹*CPHT, UMR 7644 of CNRS, Ecole Polytechnique, Palaiseau, France*

²*LPTMC, UMR 7600 of CNRS, Université P. et M. Curie, Paris, France*

³*LPMMC, UMR 5493 of CNRS, Université J. Fourier, Grenoble, France*

(Dated: July 7, 2008)

We consider the ground state of the homogeneous electron gas and we prove that a Hartree-Fock solution, motivated by previous simulations, has lower energy than the Fermi gas in the large density limit. This solution is a metallic phase : the density modulation corresponds to a partially occupied crystal (the number of sites is larger than the number of electrons).

I. INTRODUCTION

The homogeneous electron gas is one of the fundamental models in condensed matter physics. Despite its simplicity - the system consists of electrons interacting with each other through a $1/r$ potential to which a uniform positive background is added for charge neutrality - the phase diagram at zero temperature is nontrivial^{1,2,3}. In general, it is given in terms of the dimensionless parameter $r_s = (\alpha_D a_B n^{1/D})^{-1}$, where D is the space dimension, n is the electronic density, a_B the Bohr radius and α_D^D is the volume of the sphere unity. At $r_s = 0$, the ground state is the Fermi gas. At large r_s , the Hartree-Fock ground state of the electron gas is a Wigner crystal, that is a state where the charge density forms a triangular crystal with exactly one electron per lattice site. In the intermediate region the Hartree-Fock approximation is not relevant and more sophisticated methods show that the ground state of the Hamiltonian is quite different.

At small r_s , the ground state of the electron gas within Hartree-Fock is still not known⁸. Although the ground state of the ideal Fermi gas ($r_s = 0$) remains an eigenstate of the Hartree-Fock Hamiltonian of the electron gas at any density, already Wigner⁵ argued that the unpolarized Fermi gas is unstable even in the limit $r_s \rightarrow 0$. Later, Overhauser showed the instability of the unpolarized Fermi gas with respect to spin-density waves within the Hartree-Fock approximation⁶. Only recently, a Hartree-Fock study of the unpolarized three-dimensional electron gas was performed which proposes a more complicated structure of a ground state with spin-density waves⁹.

In Ref.¹⁰, we describe the results of our numerical simulations of the two-dimensional (2D), polarized electron gas at small r_s , and show that the ground state is neither a Fermi gas nor a Wigner crystal: the charge density modulation we find corresponds to a partially occupied crystal (the number of sites is larger than the number N of electrons). We refer to this solution as a metallic phase. The charge density of this metallic phase represents a triangular lattice with reciprocal generators Q_i of modulus $2k_F$. This modulation is mainly carried by the wave vectors close to the Fermi surface. In order to observe the metallic phase, the number of electrons N has to exceed a threshold ranging from $N \gtrsim 10$ at $r_s \sim 2.6$ up to $N \gtrsim 10^2$ at $r_s \sim 1$. As we shall see below, the threshold increases exponentially as r_s goes to 0 which may explain why this metallic phase has not been observed in previous simulations.

In this study, we obtain rigorous upper bounds on the energy of the metallic phase in the two-dimensional polarized case with a class of states mimicking the real states obtained numerically. These bounds are obtained in the limit $r_s \rightarrow 0$ where the calculation is simplified by the long range behavior of the interaction potential. However, our numerical simulations show that this metallic phase also exists for screened potential, but strictly speaking we cannot prove in this case the existence of such a metallic phase in the limit $r_s \rightarrow 0$. This bounds are easily extended to the unpolarized 2D case. Then we show how this proof extends to the 3D case, whereas we have no numerical simulations indicating that this phase may correspond to the actual ground state of the 3D electron gas in Hartree-Fock.

Finally, let us point out that our bounds (for instance Eq. (42)) must be considered as mathematical bounds; they only give an understanding of the behavior of the energy of the ground state, but the actual constants involved in the expressions have to be evaluated by other means.

II. NUMERICAL RESULTS

In a previous paper¹⁰, we have computed 2D Hartree-Fock states of lower energy than the Fermi gas for values of r_s about 2.

Such a state Ψ is obtained as a Slater determinant $\bigwedge_{i=1}^N \psi_i$ where N is the number of electrons and $\{\psi_i\}$ is a set of orthonormal vectors. Only the space generated by the single particle wavefunctions ψ_i 's is relevant, and in order to




FIG. 1: Numerical values of b_k for 499 electrons in two dimensions at $r_s = 2$.

understand our numerical results we need to choose a canonical representation of the ψ_i 's.

The natural choice is to use the natural base of the Fermi gas $\{\phi_i\}_{i=1\dots N}$ corresponding to some indexation of the plane waves of the Fermi sphere.

Let M be the square matrix defined by $M_{ij} = \langle \phi_i | \psi_j \rangle$. The Singular Value Decomposition (SVD) of M is $M = U\sigma V$ where U and V are unitary matrices and σ is a diagonal positive matrix.

Let us define

$$\phi'_i = \sum_j \bar{U}_{ij}^{-1} \phi_j \quad (1)$$

$$\psi'_i = \sum_j V_{ji}^{-1} \psi_j. \quad (2)$$

Then $\{\phi'_i\}$ and $\{\psi'_i\}$ are also two sets of N orthonormal vectors satisfying $\langle \phi'_i | \psi'_j \rangle = \delta_{ij} \sigma_i$.

They are the nearest bases of $\text{Span}(\{\phi_i\})$ and $\text{Span}(\{\psi_i\})$ and they define a canonical unitary operator W from $\text{Span}(\{\phi_i\})$ onto $\text{Span}(\{\psi_i\})$ by:

$$W \sum_i \lambda_i \phi'_i = \sum_i \lambda_i \psi'_i \quad (3)$$

In particular, we have:

$$W \phi_i = W \sum_j \bar{U}_{ij} \phi'_j \quad (4)$$

$$= \sum_j \bar{U}_{ij} \psi'_j \quad (5)$$

$$= \sum_j \bar{U}_{ij} \sum_k V_{kj}^{-1} \psi_k \quad (6)$$

$$= \sum_j \bar{U} \bar{V}_{ik} \psi_k \quad (7)$$

Thus the natural basis of $\text{Span}(\{\psi_i\})$ associated to the basis $\{\phi_i\}_{i=1\dots N}$ is $\{W \phi_i\}_{i=1\dots N}$.

Numerically we have chosen the ψ_i 's in this way and it appears that, if r_s is not too large, ψ_i is close to ϕ_i at least for i associated to a wave vector not too close to the Fermi surface. Thus the largest amplitude of ψ_i , in the k space, is for the k_i corresponding to ϕ_i . Fig.1 represents the next largest amplitude of ψ_i , that we denote b_{k_i} , for 499 electrons at $r_s = 2$ in 2D.

As r_s decreases, the other amplitudes of the ψ_i 's become very small. Furthermore, the wave vector k'_i corresponding to b_{k_i} satisfies $k'_i - k_i = Q_\alpha$ where the generators $\{Q_\alpha\}_{\alpha=1\dots 6}$ belong to the six-fold star of a triangular lattice.

Notice that the modulus of b_k is maximal for k on a six-fold star and close to the Fermi surface. In the direct space, the Q_α 's induce a modulation corresponding to a triangular lattice for the charge density. But while this lattice forms a Wigner crystal with one electron per site for $r_s > 2.6$, for smaller r_s it is a denser lattice corresponding to a crystal with an incomplete band filling. Indeed, at any r_s the electrostatic interaction favors a periodic distribution of charges and at small r_s only k -vectors close to Fermi surface can be modulated. The optimal solution is to choose $\|Q_\alpha\| = 2k_F$ which is larger than the reciprocal lattice vector of the Wigner case. In the following, we consider analytic solutions analogous to those of our numerical results and we prove that their energies are lower than the energies of the Fermi gas as r_s goes to 0.

III. ENERGY OF THE FERMI GAS: POLARIZED CASE

We consider the Hamiltonian of N electrons in a 2D or 3D square box of volume Ω with periodic boundary conditions.

$$H = -\frac{\hbar^2}{2m}\Delta + \frac{e^2}{2}V \quad (8)$$

where V is the 2-body Coulomb potential $\sum_{i \neq j} 1/|r_i - r_j|$, the electron mass is m , and e is its charge. It is convenient to choose Hartree as the unit of energy, $Ha = \hbar^2/(ma_B^2)$, where $a_B = \hbar^2/(me^2)$ is the Bohr radius. We get:

$$H = \frac{a_B^2}{2}(-\Delta + \frac{1}{a_B}V) \quad (9)$$

Let ψ_n be an orthonormalized set of N vectors of $L^2(\Omega)$. They define the N -particle Slater determinant $\Psi = \bigwedge_n \psi_n$. And the energy of Ψ is:

$$\mathcal{E} = \langle \Psi | H | \Psi \rangle = \frac{a_B^2}{2} \left(-\sum_n \langle \psi_n | \Delta | \psi_n \rangle + \frac{1}{a_B} \sum_{n,n'} \langle \psi_n \wedge \psi_{n'} | v | \psi_n \wedge \psi_{n'} \rangle \right) \quad (10)$$

where v is defined as:

$$\langle \varphi_1 \otimes \varphi_2 | v | \psi_1 \otimes \psi_2 \rangle = \int dx dy \overline{\varphi_1}(x) \overline{\varphi_2}(y) \frac{1}{||x - y||} \psi_1(x) \psi_2(y). \quad (11)$$

In order to avoid problems due to the Coulomb singularity, we introduce the jellium model and define the potential acting on the plane waves ϕ_k as:

$$\langle \phi_k \otimes \phi_{k'} | v | \phi_{k+q} \otimes \phi_{k'-q} \rangle = \frac{\pi}{\Omega} \left(\frac{2}{|q|} \right)^{D-1} \quad (12)$$

for $q \neq 0$ and 0 otherwise, so that the total charge of the electrons is compensated by a positive background charge. The Fermi gas is defined by $\Phi = \bigwedge_{|k| < k_F} \phi_k$ where $(\alpha_D k_F^D) = (2\pi)^D N / \Omega$ and α_D^D is the volume of the unit sphere.

$$\mathcal{E}_{FG} = \langle \Phi | H | \Phi \rangle = \frac{a_B^2}{2} \left(\sum_{|k| < k_F} k^2 - \frac{2^{D-1}\pi}{a_B \Omega} \sum_{|k|, |k'| < k_F} \frac{1}{|k - k'|^{D-1}} \right) \quad (13)$$

As Ω goes to ∞ with Ω/N fixed, the thermodynamic limit for the energy per particle is obtained by the substitution $\sum_k \rightarrow \frac{\Omega}{(2\pi)^D} \int dk$:

$$\begin{aligned} \frac{\mathcal{E}_{FG}}{N} &= \frac{a_B^2}{2} \frac{\Omega}{N(2\pi)^D} \left(\int_{|k| < k_F} dk k^2 - \frac{1}{a_B 2\pi^{D-1}} \int_{|k|, |k'| < k_F} dk dk' \frac{1}{|k - k'|^{D-1}} \right) \\ &= \frac{a_B^2}{2} \frac{\Omega}{N(2\pi)^D} k_F^{D+2} \left(\int_{|k| < 1} dk k^2 - \frac{1}{a_B k_F 2\pi^{D-1}} \int_{|k|, |k'| < 1} dk dk' \frac{1}{|k - k'|^{D-1}} \right) \end{aligned} \quad (14)$$




FIG. 2: The circle is the Fermi surface. The shaded surfaces are the regions where $b(k)$ is nonzero. The new state ψ_k mixing ϕ_k and ϕ_{k+Q_k} is now resonant with $\psi_{k'}$.

From the definition of $r_s = (\alpha_D a_B n^{1/D})^{-1}$ and k_F , it follows that $k_F \alpha_D^2 r_s a_B = 2\pi$. Thus, we have:

$$\frac{\mathcal{E}_{FG}}{N} = \frac{2\pi^2}{\alpha_D^{D+4} r_s^2} \left(\int_{|k|<1} dk k^2 - \frac{r_s \alpha_D^2}{4\pi^D} \int_{|k|,|k'|<1} dk dk' \frac{1}{|k - k'|^{D-1}} \right) \quad (15)$$

which gives for $D = 2$ ($\alpha_2^2 = \pi$):

$$\frac{\mathcal{E}_{FG}}{N} = \frac{2}{\pi r_s^2} \left(\int_{|k|<1} dk k^2 - \frac{r_s}{4\pi} \int_{|k|,|k'|<1} dk dk' \frac{1}{|k - k'|} \right) \quad (16)$$

IV. HARTREE-FOCK UPPER BOUNDS: POLARIZED 2D CASE

We restrict ourself to the 2D polarized case and we want to estimate the energy for a class of states inspired by our numerical results. Let us consider a state $\Psi = \bigwedge_{|k|<k_F} \psi_k$ where:

$$\psi_k = a_k \phi_k + b_k \phi_{k+Q_k}$$

with Q_k in $\{-2k_F(\cos p\pi/3, \sin p\pi/3)\}_{p=0\dots 5}$. For $k = |k|(\cos \theta, \sin \theta)$ we choose Q_k such that $|k + Q_k|$ is minimal; that is, we choose p as the integer part of $(3\theta/\pi + 1/2)$ and we must assume b_k is zero if k is zero or $\theta = \pi/6 + n\pi/3$. Furthermore, we assume that a_k and b_k are real positive number and invariant thru the rotation of $2n\pi/6$ and the symmetry $\theta \rightarrow -\theta$ (i.e. the dihedral group D_6). The ψ_k 's are normalized, so that $a_k^2 + b_k^2 = 1$ and $b_k = 0$ if $|k \cdot Q_k| < 2k_F^2(1 - \epsilon)$ (i.e. b_k is not zero only in the vicinity of $\{k_F(\cos p\pi/3, \sin p\pi/3)\}_{p=0\dots 5}$), see Fig.2.

Thus the limit energy per particle is given by:

$$\frac{\mathcal{E}}{N} = \frac{2}{\pi r_s^2} \left(\int_{|k|<1} dk \langle \psi_k | -\Delta | \psi_k \rangle + \frac{r_s}{4\pi} \int_{|k|,|k'|<1} dk dk' \frac{\Omega}{2\pi} \langle \psi_k \wedge \psi_{k'} | v | \psi_k \wedge \psi_{k'} \rangle \right) \quad (17)$$

where, as in (16), the k 's have been renormalized by k_F and thus $|Q_k| = 2$.

We define ΔE by:

$$\frac{\mathcal{E} - \mathcal{E}_{FG}}{N} = \frac{2}{\pi r_s^2} \Delta E \quad (18)$$

Then

$$\Delta E = \int_{|k|<1} dk [\langle \psi_k | -\Delta | \psi_k \rangle - k^2] + \frac{r_s}{4\pi} \Delta E_V \quad (19)$$

where

$$\Delta E_V = \int_{|k|, |k'| < 1} dk dk' \left(\frac{\Omega}{2\pi} \langle \psi_k \wedge \psi_{k'} | v | \psi_k \wedge \psi_{k'} \rangle + \frac{1}{|k - k'|} \right) \quad (20)$$

A. Potential energy contribution: ΔE_V

Setting $v_q = 1/|q|$:

$$\begin{aligned} \frac{\Omega}{2\pi} \langle \psi_k \wedge \psi_{k'} | v | \psi_k \wedge \psi_{k'} \rangle + v_{k-k'} &= (v_{k-k'} - v_{k-k'-Q_{k'}}) b_{k'}^2 a_k^2 \\ &+ (v_{k-k'} - v_{k+Q_k-k'}) b_k^2 a_{k'}^2 \\ &+ (v_{k-k'} - v_{k+Q_k-Q_{k'}-k'}) b_k^2 b_{k'}^2 \\ &+ 2v_{Q_k} a_k b_k a_{k'} b_{k'} (\delta_{Q_k+Q_{k'}} + \delta_{Q_k-Q_{k'}}) \\ &- 2v_{k-k'} a_k b_k a_{k'} b_{k'} \delta_{Q_k-Q_{k'}} \\ &- (v_{k+Q_k-k'} + v_{k-k'-Q_{k'}}) a_k a_{k'} b_k b_{k'} \delta_{Q_k+Q_{k'}} \end{aligned} \quad (21)$$

Eq.(20), may be divided into 4 parts:

- $\{b_{k'} = 0, b_k = 0\}$: the contribution is zero.
- $\{b_{k'} = 0, b_k \neq 0\}, \{b_{k'} \neq 0, b_k = 0\}$: both cases are equivalent.

For $\{b_{k'} = 0, b_k \neq 0\}$, the integrant of Eq. 20 is:

$$\frac{\Omega}{2\pi} (\psi_k \wedge \phi_{k'}, v \psi_k \wedge \phi_{k'}) + \frac{1}{|k - k'|} = (v_{k-k'} - v_{k+Q_k-k'}) b_k^2 \quad (22)$$

Let S_0 be the sector of unit disk between $-\pi/6$ and $\pi/6$ (see Fig. 2); then in this sector $Q_k = (-2, 0)$ and by symmetry:

$$\int_{b_{k'}=0} dk dk' (v_{k-k'} - v_{k+Q_k-k'}) b_k^2 = 6 \int_{k \in S_0, b_{k'}=0} dk dk' (v_{k-k'} - v_{k+Q_k-k'}) b_k^2 \quad (23)$$

$$= 6 \int_{k \in S_0, b_{k'}=0} dk dk' (v_{k-k'} - v_{\tilde{k}-k'}) b_k^2 \quad (24)$$

$$\leq C\epsilon^3 + 6 \int_{k \in S_0, |k'_x| < 1-\epsilon} dk dk' (v_{k-k'} - v_{\tilde{k}-k'}) b_k^2 \quad (25)$$

where $\tilde{k} = (2 - k_x, k_y)$. In S_0 , $k = (k_x, k_y)$ where k_x is close to 1 and setting $k_x = 1 - x$, we assume from now that $b_k = b(x/\epsilon)$.

In Appendix A we prove that:

$$\int_{k \in S_0, |k'_x| < 1-\epsilon} dk dk' (v_{k-k'} - v_{k+Q_k-k'}) b_k^2 \leq 8\epsilon^2 \sqrt{2\epsilon} [\ln \epsilon^{-1} + O(1)] \int_0^1 dx b^2(x) x \sqrt{x} \quad (26)$$

- $\{b_{k'} \neq 0, b_k \neq 0\}$:

By symmetry we can assume that k belongs to S_0 . If $k' \notin S_0 \cup S_3$ all the v appearing in (21) are uniformly bounded. And since the k -volume for each sector goes like $\epsilon\sqrt{\epsilon}$, the contribution of these terms is bounded by $C\epsilon^3$. In the same way $v_{k-k'}$ is bounded when $k' \in S_3$ and $v_{k+Q_k-k'}$ is bounded when $k' \in S_0$.

Thus setting:

$$f := a_k^2 b_{k'}^2 + b_k^2 a_{k'}^2 - 2a_k a_{k'} b_k b_{k'} = (a_k b_{k'} - b_k a_{k'})^2 \quad (27)$$

$$g := a_k^2 b_{k'}^2 + b_k^2 a_{k'}^2 + 2a_k a_{k'} b_k b_{k'} = (a_k b_{k'} + b_k a_{k'})^2 \quad (28)$$

one can check that:

$$\int_{b_k, b_{k'} \neq 0} \frac{\Omega}{2\pi} \langle \psi_k \wedge \psi_{k'} | v | \psi_k \wedge \psi_{k'} \rangle + v_{k-k'} \leq C\epsilon^3 + 6 \int_{k, k' \in S_0} dk dk' (v_{k-k'} f - v_{k+Q_k+k'} g) \quad (29)$$

In Appendix B we prove that

$$\int_{k, k' \in S_0} dk dk' (v_{k-k'} f - v_{k+Q_k+k'} g) \leq 4\epsilon^2 \sqrt{2\epsilon} [\ln \epsilon^{-1} + O(1)] \int_0^1 dx \sqrt{x} \int_x^1 dx' (f(\epsilon x, \epsilon x') - g(\epsilon x, \epsilon x')) \quad (30)$$

Thus, summing the four contribution gives:

$$\Delta E_V \leq C\epsilon^3 + 6\epsilon^2 \sqrt{2\epsilon} [\ln \epsilon^{-1} + O(1)] \int_0^1 dx \sqrt{x} \left(16b^2(x)x + 4 \int_x^1 dx' (f(x, x') - g(x, x')) \right) \quad (31)$$

B. Kinetic energy contribution:

The variation of the kinetic energy is given by:

$$\int_{|k| < 1} dk [\langle \psi_k | -\Delta | \psi_k \rangle - k^2] = 6 \int_{k \in S_0} dk (\langle \psi_k | -\Delta | \psi_k \rangle - k^2) \quad (32)$$

$$= 6 \int_0^\epsilon dx \, 2y_m \, 4xb^2(x/\epsilon) \quad (33)$$

$$\leq 6 \times 8\epsilon^2 \sqrt{2\epsilon} \int_0^1 dx \sqrt{x} \, xb^2(x) \quad (34)$$

C. Total energy:

Inserting Eqs.(31,34) in Eq. (19), the variation of the total energy from the Fermi gas energy becomes:

$$\begin{aligned} \Delta E &\leq 6\epsilon^2 \sqrt{2\epsilon} \int_0^1 dx \sqrt{x} \left(8xb^2(x) + \frac{r_s}{4\pi} [\ln \epsilon^{-1} + O(1)] \left(16b^2(x)x + 4 \int_x^1 dx' (f(x, x') - g(x, x')) \right) \right) \\ &= 6 \times 8\epsilon^2 \sqrt{2\epsilon} \int_0^1 dx \sqrt{x} \left(xb^2(x) + \frac{r_s}{2\pi} [\ln \epsilon^{-1} + O(1)] \left(b^2(x)x - a(x)b(x) \int_x^1 dx' a(x')b(x') \right) \right) \end{aligned} \quad (35)$$

Let us set

$$\delta = \epsilon^2 \sqrt{\epsilon} \quad (36)$$

$$I_1 = \int_0^1 dx \sqrt{x} xb^2(x) \quad (37)$$

$$I_2 = \frac{1}{5\pi} \int_0^1 dx \sqrt{x} \left(-b^2(x)x + a(x)b(x) \int_x^1 dx' a(x')b(x') \right) \quad (38)$$

Then

$$\Delta E \leq 6 \times 8\sqrt{2}\delta [I_1 - r_s I_2 (\ln \delta^{-1} + O(1))] \quad (39)$$

If $I_2 > 0$, as r_s goes to 0, ΔE is minimal in Eq. (39) for δ defined by:

$$\delta_{\min} = \frac{1}{e} \exp \left(-\frac{I_1}{I_2 r_s} \right) \quad (40)$$

and finally inserting $\delta_{\min}$ in Eq. (39) gives:

$$\Delta E \lesssim -\frac{6 \times 8\sqrt{2}}{e} \exp \left(-\frac{I_1}{I_2 r_s} \right) r_s I_2 \quad (41)$$

We now have to find a solution $b(x)$ such that I_2 is positive. Choosing $b(x) = b_0$ or $b(x) = b_0(1 - x)$ leads to negative I_2 . In the Appendix C, as r_s goes to 0 we find a family of b leading to :

$$\Delta E \lesssim -r_s \exp\left(-\frac{5\pi}{3r_s} + \frac{O(1)}{\sqrt{r_s}}\right) \quad (42)$$

Though such a bound is correct in the thermodynamic limit, this behavior is not so relevant for finite systems. Indeed, numerical systems consider about 10^3 electrons and, as we shall see later, the Fermi gas becomes then the ground state for $r_s \lesssim 1$. Thus the asymptotic bound (42) is not very helpful in the real world. Nevertheless, for realistic r_s , one can choose a suitable function b and evaluate numerically I_1 and I_2 . For instance, with $b = b_\eta$ as in (88) of Appendix C and $\eta = 0.001$ we get

$$\Delta E \lesssim -2.6 \times 10^{-4} r_s \exp\left(-\frac{18.5}{r_s}\right) \quad (43)$$

V. HF UPPER BOUNDS: UNPOLARIZED 2D CASE

We consider now the spin of the electrons, and we restrict to the case where half the electrons have a spin up and the others have a spin down.

We can choose a solution as the product of 2 Slaters Ψ^+ and Ψ^- for the spins up and down.

$$\frac{2}{a_B^2} \mathcal{E}^{NP} = - \sum_n \langle \psi_n^+ | \Delta | \psi_n^+ \rangle + \frac{1}{a_B} \sum_{n,n'} \langle \psi_n^+ \wedge \psi_{n'}^+ | v | \psi_n^+ \wedge \psi_{n'}^+ \rangle \quad (44)$$

$$- \sum_n \langle \psi_n^- | \Delta | \psi_n^- \rangle + \frac{1}{a_B} \sum_{n,n'} \langle \psi_n^- \wedge \psi_{n'}^- | v | \psi_n^- \wedge \psi_{n'}^- \rangle \quad (45)$$

$$+ \frac{1}{a_B} \left(\sum_{n,n'} \langle \psi_n^+ \otimes \psi_{n'}^- | v | \psi_n^+ \otimes \psi_{n'}^- \rangle + \sum_{n,n'} \langle \psi_n^- \otimes \psi_{n'}^+ | v | \psi_n^- \otimes \psi_{n'}^+ \rangle \right) \quad (46)$$

The first terms (44) and (45) can be evaluated as in (18) noticing that the definition of k_F becomes $k_F^2 = 4\pi N_\pm / \Omega = 2\pi N / \Omega$. So the r_s in the evaluation of ΔE is now $r_s \sqrt{2}$ and:

$$\frac{\mathcal{E}^{NP} - \mathcal{E}_{FG}^{NP}}{N} = \frac{1}{\pi r_s^2} \Delta E(r_s \sqrt{2}) + (46) \quad (47)$$

The extra terms (46) coming from the direct potential provide contributions like $a_n^+ b_n^+ a_n^- b_n^-$, times a positive factor. These contributions are regular as r_s goes to 0, and does not modify the asymptotic energy.

Nevertheless, these contributions depends on the signs of $b_n^\pm$ in the two Slaters Ψ^+ and Ψ^- . Thus, since the energy of one Slater does not depend on the global sign of the $b_n^\pm$'s, the minimal state is obtained by choosing $b_n^- = -b_n^+$.

For such a state, the main difference with the polarized case is that the charge density becomes flat, though the spin density is not.

Finally, we can compare our solutions with previous solutions proposed by Fedders and Martin⁷. They consider a more complicated modulation of the states near the center of Fermi sphere. While their results may be correct, they missed the dominant contribution coming from the states near the Fermi surface. Indeed, our computations give nonzero contribution for the states near the center, but for $r_s = 1.5$ b_k is about 10^{-5} and the energy benefit, following their formula, must be of order 10^{-20} of the energy benefit of the external states.

VI. THE METALLIC PHASE IN FINITE SYSTEMS

For finite systems of N electrons, the minimum requirement is that the surface $|k \cdot Q_k| > 2k_F^2(1 - \epsilon)$, so that it contains at least one plane wave of the finite, discrete system. This gives the condition $N\epsilon\sqrt{\epsilon} > 1$ and from Eqs. (36,40), this leads to:

$$N > \exp\left(\frac{3I_1}{5I_2 r_s}\right) \quad (48)$$

With our approximate behavior for I_2/I_1 in two dimensions, we find:

$$N > \exp\left(\frac{3\pi}{r_s}\right) \quad (49)$$

i.e. $N > 500$ for $r_s = 1.8$. This bound is compatible with our numerical simulations where the metallic phase disappears at $rs = 1$ for $N = 500$. Furthermore, this may explain why this metallic phase has not been observed in previous simulations.

VII. HARTREE-FOCK UPPER BOUNDS: POLARIZED 3D CASE

Mutatis mutandis, the 3D case works in the same way.

The potential term v_k is now $1/\|k\|^2$ but the dominant terms are provide now from integrals like:

$$I(k_x, k_{x'}) = \int_{\|k\|, \|k'\| \leq 1} dk_y dk_z dk'_y dk'_z \frac{1}{\|k - k'\|^2} \quad (50)$$

where k_x is close to k'_x .

Let us suppose that $k_x > k'_x$, then

$$I(k_x, k_{x'}) = \int_0^{\sqrt{2-k_x^2-k_x'^2}} \mu(du) \frac{1}{(k_x - k'_x)^2 + u^2} \quad (51)$$

where u stands for $((k_y - k'_y)^2 + (k_z - k'_z)^2)^{1/2}$ and $\mu(du)$ stands for the distribution of u . Here we are interested in small values of u . One can check that $\mu(u, u + \epsilon) \approx 2\pi^2(1 - k_x^2)\epsilon u$ provided that u is sufficiently small ((k_y, k_z) may be any point in disk of radius $\sqrt{1 - k_x^2}$ and (k'_y, k'_z) may belong to a thin ring around (k_y, k_z))

$$I(k_x, k_{x'}) \approx \pi^2(1 - k_x^2) \int_0^{2\sqrt{2\epsilon}} 2udu \frac{1}{(k_x - k'_x)^2 + u^2} \quad (52)$$

$$\approx -\pi^2(1 - k_x^2) \ln(k_x - k'_x)^2 \quad (53)$$

This must be compared with the 2D case $I(k_x, k_{x'}) \approx -2\sqrt{1 - k_x^2} \ln(k_x - k'_x)^2$. Mutatis mutandis, (Eq. 19) becomes:

$$\Delta E = \int_{|k| < 1} dk \left(\langle \psi_k | - \Delta | \psi_k \rangle - k^2 \right) + \kappa r_s \Delta E_V \quad (54)$$

where $\kappa = \alpha_3^2/(4\pi^3)^2$ and $\alpha_3^3 = 4\pi/3$. Now, in ΔE_V the $\sqrt{2\epsilon x}$ factors have to be replaced by $\pi^2 \epsilon x$:

$$\Delta E_V \leq C\epsilon^4 + 16z\pi^2\epsilon^3 [\ln \epsilon^{-1} + O(1)] \int_0^1 dx \, x \left(b^2(x)x - a(x)b(x) \int_x^1 dx' a(x')b(x') \right) \quad (55)$$

where z is the number of sectors of the Fermi sphere. The variation of the kinetic energy is now

$$\int_{|k| < 1} dk \left(\langle \psi_k | - \Delta | \psi_k \rangle - k^2 \right) = z8\pi\epsilon^3 \int_0^1 dx \, x^2 b^2(x) \quad (56)$$

Thus setting

$$I_1 = \int_0^1 dx \, x^2 x b^2(x) \quad (57)$$

$$I_2 = \frac{2\kappa\pi}{3} \int_0^1 dx \, x \left(-b^2(x)x + a(x)b(x) \int_x^1 dx' a(x')b(x') \right) \quad (58)$$

$$\Delta E \leq z8\pi\epsilon^3 (I_1 - 3r_s I_2 [\ln \epsilon^{-1} + O(1)])$$

Once more, if we find a b such that I_2 is positive, we have:

$$\Delta E \lesssim -\frac{z8\pi}{e} \exp\left(-\frac{I_1}{I_2 r_s}\right) r_s I_2 \quad (59)$$

The operator A of Appendix C is now modified. But the same analysis show that the spectrum of A is still $(0, 4)$, so that the asymptotic bound for small r_s is:

$$\Delta E \lesssim -\frac{z8\pi}{e} \exp\left(-\frac{1}{6\kappa\pi r_s} + o(1/r_s)\right) r_s \quad (60)$$

VIII. CONCLUSION

We have proven upper bounds for the energy of a metallic state below the Fermi gas energy. To our knowledge, it is the first time that rigorous upper bounds for the ground state energy of the polarized electron gas are obtained going beyond the simple Fermi gas states. The main interest of this proof is that it focuses on the small r_s region where the Hartree-Fock approximation is relevant and that the upper bounds were obtained using metallic states which correspond to solutions obtained by numerical simulations.

In order to judge their relevance for the true ground state of the electron gas in the high density region, we have to consider correlation effects beyond the Hartree-Fock approximation. Due to the long-range, singular behavior of the Coulomb-potential, the perturbation expansion has to be rearranged which amounts to an effective screening of the electron interaction. Unfortunately a rigorous extension of our proof to the case of a screened Coulomb potential is not straightforward. However, numerical calculations provide evidence for the stability of the metallic states in the 2D polarized system within Hartree-Fock, so that these states are relevant candidates for the true ground state of the two-dimensional electron gas.

IX. APPENDIX A

We have to estimate

$$I(f) = \int_{\substack{\|k\|, \|k'\| \leq 1 \\ 1-k_x < \epsilon, |k'_x| < 1-\epsilon}} dk dk' (v_{k-k'} - v_{\tilde{k}-k'}) f(1-k_x) \quad (61)$$

where $\tilde{k} = (2-k_x, k_y)$ and f is a positive function.

$$\int dk' (v_{k-k'} - v_{\tilde{k}-k'}) = \int dk'_x \operatorname{asinh} \frac{y'_m - k_y}{k_x - k'_x} + \operatorname{asinh} \frac{y'_m + k_y}{k_x - k'_x} - \operatorname{asinh} \frac{y'_m - k_y}{2-k_x - k'_x} - \operatorname{asinh} \frac{y'_m + k_y}{2-k_x - k'_x} \quad (62)$$

where $y'_m = \sqrt{1-k'^2_x}$. And since $\operatorname{asinh} x - \operatorname{asinh} y \leq \ln x/y$ for $x > y > 0$:

$$\int dk' (v_{k-k'} - v_{\tilde{k}-k'}) \leq \int_{-1+\epsilon}^{1-\epsilon} dk'_x 2 \ln \frac{2-k_x - k'_x}{k_x - k'_x} \quad (63)$$

We set $k_x = 1-x$ and $y_m = \sqrt{2x-x^2}$,

$$I(f) \leq \int_0^\epsilon dx f(x) 2y_m \int_{-1+\epsilon}^{1-\epsilon} dk'_x 2 \ln \frac{1+x-k'_x}{1-x-k'_x} \quad (64)$$

$$= 4 \int_0^\epsilon dx f(x) y_m \int_\epsilon^{2-\epsilon} du \ln \frac{u+x}{u-x} \quad (65)$$

$$\leq 4 \int_0^\epsilon dx f(x) y_m \int_\epsilon^{2-\epsilon} du \frac{2x}{u-x} \quad (66)$$

$$= 8\epsilon^2 \sqrt{2\epsilon} [\ln \epsilon^{-1} + O(1)] \int_0^1 dx f(\epsilon x) x \sqrt{x} \quad (67)$$

X. APPENDIX B

We have to estimate

$$I(f, g) = \int_{\substack{\|k\|, \|k'\| \leq 1 \\ 1-k_x, 1-k'_x < \epsilon}} dk dk' (v_{k-k'} f - v_{\tilde{k}+k'} g) \quad (68)$$

where $\tilde{k} = (k_x - 2, k_y)$ and f and g are positive functions of $1 - k_x$ and $1 - k_{x'}$.

Setting $k_x = 1 - x$, $k'_x = 1 - x'$, $k_y = y$, $k'_y = y'$, and $r_{\pm} = \sqrt{(x \pm x')^2 + (y - y')^2}$, Eq.68 can be rewritten:

$$\begin{aligned} I(f, g) &= \int_0^\epsilon dx \int_0^\epsilon dx' \int dy dy' \left(\frac{1}{r_-} f - \frac{1}{r_+} g \right) \\ &= 2 \int_0^\epsilon dx \int_x^\epsilon dx' \int dy dy' \left(\frac{1}{r_-} f - \frac{1}{r_+} g \right) \end{aligned} \quad (69)$$

where y and y' must satisfy $(1 - x)^2 + y^2 \leq 1$ and $(1 - x')^2 + y'^2 \leq 1$.

Since $\operatorname{asinh} x \leq \ln 2(x + 1)$, the first term in 69 is bounded by:

$$2 \int_0^\epsilon dx \int_x^\epsilon dx' \int dy dy' \frac{1}{r_-} f = 2 \int_0^\epsilon dx \int_x^\epsilon dx' f \int_{-y_m}^{y_m} dy (\operatorname{asinh} \frac{y'_m + y}{x' - x} + \operatorname{asinh} \frac{y'_m - y}{x' - x}) \quad (70)$$

$$\leq 2 \int_0^\epsilon dx \int_x^\epsilon dx' f \int_{-y_m}^{y_m} dy 2 \operatorname{asinh} \frac{2y'_m}{x' - x} \quad (71)$$

$$\leq 4 \int_0^\epsilon dx \int_x^\epsilon dx' f 2y_m \ln(2 + \frac{4y'_m}{x' - x}) \quad (72)$$

$$\leq 4\epsilon^2 \sqrt{2\epsilon} [\ln(\epsilon^{-1}) + O(1)] \int_0^1 dx \int_x^1 dx' f(\epsilon x, \epsilon x') \sqrt{x} \quad (73)$$

On the other hand, using $\operatorname{asinh} x \geq \ln 2x$, the last term of (69) is:

$$2 \int_0^\epsilon dx \int_x^\epsilon dx' g \int dy dy' \frac{1}{r_+} = 2 \int_0^\epsilon dx \int_x^\epsilon dx' g \int_{-y_m}^{y_m} dy \operatorname{asinh} \frac{y'_m - y}{x + x'} + \operatorname{asinh} \frac{y'_m + y}{x + x'} \quad (74)$$

$$\geq 2 \int_0^\epsilon dx \int_x^\epsilon dx' g \int_{-y_m}^{y_m} dy \ln 4 \frac{y_m'^2 - y^2}{(x + x')^2} \quad (75)$$

$$\geq 4 \int_0^\epsilon dx \int_x^\epsilon dx' g y_m [\ln \epsilon^{-1} + O(1)] \quad (76)$$

$$\geq 4\epsilon^2 \sqrt{2\epsilon} [\ln \epsilon^{-1} + O(1)] \int_0^1 dx \int_x^1 dx' g(\epsilon x, \epsilon x') \sqrt{x} \quad (77)$$

And we have:

$$I(f, g) \leq 4\epsilon^2 \sqrt{2\epsilon} [\ln \epsilon^{-1} + O(1)] \int_0^1 dx \sqrt{x} \int_x^1 dx' (f(\epsilon x, \epsilon x') - g(\epsilon x, \epsilon x')) \quad (78)$$

XI. APPENDIX C

Here we provide exact bounds on I_1 and I_2 given by (37, 57).

In order to estimate I_2 we introduce the linear operator A :

$$Af(x) = \frac{1}{2x} \int_x^1 f(y) dy + \frac{1}{2x\sqrt{x}} \int_0^x f(y) \sqrt{y} dy \quad (79)$$

defined on the Hilbert space of the functions on $[0, 1]$ with the scalar product:

$$\langle f | g \rangle = \int_0^1 x \sqrt{x} \overline{f(x)} g(x) dx \quad (80)$$

Then A is a bounded symmetric operator and:

$$I_2/I_1 = \frac{1}{5\pi} \left(\frac{\langle ab|Aab\rangle}{\|b\|^2} - 1 \right) \quad (81)$$

The unitary operator $f(x) \rightarrow g(y) = f(e^{-y})e^{-5/4y}$ from $L^2([0, 1], x\sqrt{x}dx)$ onto $L^2([0, +\infty], dx)$ maps the operator A onto the operator $\tilde{A}$:

$$\tilde{A}g(x) = \frac{e^{-x/4}}{2} \int_0^x e^{y/4} g(y) dy + \frac{e^{x/4}}{2} \int_x^{+\infty} e^{-y/4} g(y) dy \quad (82)$$

Then

$$\tilde{A}e^{ikx} = \frac{1}{4(1/16 + k^2)} e^{ikx} - \frac{1}{1/2 + i2k} e^{-x/4} \quad (83)$$

Thus setting:

$$g_k(x) = \frac{1}{|1 + i4k|} [(1 + i4k)e^{ikx} - (1 - i4k)e^{-ikx}] \quad (84)$$

$\{g_k\}_{k>0}$ is a full set of pseudo-eigenvectors satisfying:

$$\tilde{A}g_k = \frac{1}{4(1/16 + k^2)} g_k$$

Thus the spectrum of $\tilde{A}$ is $(0, 4)$ and the spectral measure is purely absolutely continuous; the largest spectral value is 4 with a pseudo-eigenvector $g_4(x) = x + 4$ corresponding to $f_4(x) = x^{-5/4}(4 - \ln x)$.

But $\|f_4\|$ is infinite and f_4 diverges at 0. The next step is to choose a family of functions b_η such that $a_\eta = \sqrt{1 - b_\eta}$ is defined and $\langle a_\eta b_\eta | A a_\eta b_\eta \rangle / \|b_\eta\|^2$ is close to 4.

Thus setting $f_\eta(x) = \min(f_4(x), f_4(\eta))$ for $0 < \eta \ll 1$, we have:

$$\|f_\eta\|^2 = -\frac{1}{3} \left[\ln^3 \eta - \frac{66}{5} \ln^2 \eta + O(\ln \eta) \right] \quad (85)$$

$$\langle f_\eta | A f_\eta \rangle = -\frac{4}{3} \left[\ln^3 \eta - \frac{41}{5} \ln^2 \eta + O(\ln \eta) \right] \quad (86)$$

Then:

$$\frac{\langle f_\eta | A f_\eta \rangle}{\|f_\eta\|^2} = 4 - \frac{20}{|\ln \eta|} + O(\ln^{-2} \eta) \quad (87)$$

Thus f_η is a good candidate for the linear part of the problem. Now, by the simple scaling:

$$b_\eta(x) = \frac{f_\eta(x)}{\sqrt{2}f_\eta(\eta)} \quad (88)$$

we get the nonlinear candidate satisfying $b_\eta(x) \leq 1/\sqrt{2}$, $a_\eta = \sqrt{1 - b_\eta^2}$ is well defined, $a_\eta(x) \geq 1/\sqrt{2}$ and b_η satisfies (87).

We must now estimate the simultaneous convergence of I_2/I_1 (81) and I_2 as η decreases.

$$\begin{aligned} \langle b_\eta | A b_\eta \rangle - \langle b_\eta a_\eta | A b_\eta a_\eta \rangle &= -\langle b_\eta - b_\eta a_\eta | A | b_\eta - b_\eta a_\eta \rangle + 2 \langle b_\eta - b_\eta a_\eta | A | b_\eta \rangle \\ &\leq 2 \langle b_\eta - b_\eta a_\eta | A | b_\eta \rangle \\ &= 8 \langle b_\eta - b_\eta a_\eta | b_\eta \rangle + 2 \langle b_\eta - b_\eta a_\eta | (A - 4) b_\eta \rangle \\ &\leq 8 \langle b_\eta - b_\eta a_\eta | b_\eta \rangle + 2 \|b_\eta - b_\eta a_\eta\| \|(A - 4) b_\eta\| \\ &\leq 8 \langle b_\eta - b_\eta a_\eta | b_\eta \rangle + 8 \|b_\eta - b_\eta a_\eta\| \sqrt{\langle b_\eta | (A - 4) b_\eta \rangle} \end{aligned}$$

where:

$$\begin{aligned}\langle b_\eta - b_\eta a_\eta | b_\eta \rangle &= \int_0^1 b_\eta(x)^2 [1 - a_\eta(x)] x \sqrt{x} dx \\ &\leq \int_0^1 b_\eta(x)^2 [1 - a_\eta(x)]^2 x \sqrt{x} dx \sup \frac{1}{1 - a_\eta} \\ &\leq \|b_\eta - b_\eta a_\eta\|^2 \frac{\sqrt{2}}{\sqrt{2} - 1}\end{aligned}$$

and since $b_\eta - b_\eta a_\eta > 0$

$$\begin{aligned}\|b_\eta - b_\eta a_\eta\|^2 &= \|b_\eta\|^2 - \|b_\eta a_\eta\|^2 - 2 \langle b_\eta - b_\eta a_\eta | b_\eta a_\eta \rangle \\ &\leq \|b_\eta\|^2 - \|b_\eta a_\eta\|^2 \\ &= \|b_\eta^2\|^2\end{aligned}$$

By direct computation:

$$\|b_\eta^2\|^2 \leq \|b_\eta\|^2 \frac{6}{5|\ln \eta|}$$

for η small enough, and thus:

$$\frac{\langle b_\eta | A b_\eta \rangle}{\|b_\eta\|^2} - \frac{\langle b_\eta a_\eta | A b_\eta a_\eta \rangle}{\|b_\eta\|^2} \leq \frac{8}{|\ln \eta|} \left(\frac{\sqrt{2}}{\sqrt{2} - 1} \frac{6}{5} + 2\sqrt{6} \right) \quad (89)$$

And finally, from (87) for b_η and (89) and $I_1 = \|b_\eta\|^2$, (81) gives:

$$I_2/I_1 \geq \frac{1}{5\pi} \left(3 - \frac{C}{|\ln \eta|} \right) + O(\ln^{-2} \eta) \quad (90)$$

$$I_2 \leq \frac{2}{15\pi} \eta^{5/2} |\ln^3 \eta| \left[1 + O\left(\frac{1}{|\ln \eta|} \right) \right] \quad (91)$$

where

$$C = 20 + 8 \left(\frac{\sqrt{2}}{\sqrt{2} - 1} \frac{6}{5} + 2\sqrt{6} \right) \approx 92 \quad (92)$$

Choosing η sufficiently small, this proves that ΔE is strictly negative for any $r_s > 0$; furthermore choosing η to minimize ΔE (Eq. 41), i.e. $9r_s |\ln \eta|^2 = 2C\pi$, we obtain as r_s goes to 0:

$$\Delta E \lesssim -r_s \exp \left(-\frac{5\pi}{3r_s} + \frac{O(1)}{\sqrt{r_s}} \right) \quad (93)$$

-
- ¹ B. Tanatar and D. M. Ceperley, Phys. Rev. **B 39**, 5005 (1989); C. Attaccalite, S. Moroni, P. Gori-Giorgi, and G. B. Bachelet, Phys. Rev. Lett. **88**, 256601 (2002).
² B. Bernu, L. Cândido, and D. M. Ceperley, Phys. Rev. Lett. **86**, 870 (2001).
³ H. Falakshahi and X. Waintal, Phys. Rev. Lett. **94**, 046801 (2005), X. Waintal, Phys. Rev. **B 73**, 075417 (2006).
⁴ J. R. Trail, M. D. Towler, and R. J. Needs, Phys. Rev. **B 68**, 045107 (2003).
⁵ E. P. Wigner, Trans. Faraday Soc. **34**, 678 (1938); Phys. Rev. **46**, 1002 (1934).
⁶ A. W. Overhauser, Phys. Rev. Lett. **4**, 462 (1960); Phys. Rev. **128**, 1437 (1962).
⁷ A. Fedders and P. C. Martin, Phys. Rev. **143**, 245-259 (1966).
⁸ G. F. Giuliani and G. Vignale, Quantum Theory of the Electron Liquid, Cambridge University Press, Cambridge (2005).
⁹ S. Zhang and D. M. Ceperley, Phys. Rev. Lett. **100**, 236404 (2008), arXiv:0712.1194 (2007).
¹⁰ B. Bernu, F. Delyon, M. Duneau, and M. Holzmann, cond-mat. arXiv:0804.1025